

Asymptotic analysis of the Krawtchouk polynomials by the WKB method

Diego Dominici *

Department of Mathematics
State University of New York at New Paltz
75 S. Manheim Blvd. Suite 9
New Paltz, NY 12561-2443
USA

February 1, 2008

Abstract

We analyze the Krawtchouk polynomials $K_n(x, N, p, q)$ asymptotically. We use singular perturbation methods to analyze them for $N \rightarrow \infty$, with appropriate scalings of the two variables x and n . In particular, the WKB method and asymptotic matching are used. We obtain asymptotic approximations valid in the whole domain $[0, N] \times [0, N]$, involving some special functions. We give numerical examples showing the accuracy of our formulas.

1 Introduction

Definition 1 *Let $n, N \geq 0$ be integers. The Krawtchouk polynomials $K_n(x)$ are defined by [36],*

$$K_n(x) = \sum_{k=0}^n \binom{x}{k} \binom{N-x}{n-k} q^k (-p)^{n-k} \quad (1)$$

where

$$0 < p, q < 1 \quad p + q = 1. \quad (2)$$

The binary Krawtchouk polynomials are the special case with $p = 1/2 = q$ [15], [18].

*e-mail: dominicd@newpaltz.edu

The Krawtchouk polynomials [27] are one of the families of classical orthogonal polynomials of a discrete variable [14]. They satisfy the orthogonality relation

$$\sum_{k=0}^N K_i(k) K_j(k) \varrho(k) = \binom{N}{j} (pq)^j \delta_{ij}, \quad i, j = 0, \dots, N$$

with weight function

$$\varrho(x) = \binom{N}{x} p^x q^{N-x}.$$

Writing the Krawtchouk polynomials in the extended form $K_n(x) = K_n(x, N, p, q)$, we have the symmetry formula

$$K_n(x, N, p, q) = (-1)^n K_n(N - x, N, q, p). \quad (3)$$

They also satisfy the three-term recurrence

$$(n+1)K_{n+1}(x) + pq(N-n+1)K_{n-1}(x) + [p(N-n) + nq-x]K_n(x) = 0$$

which is the main object of our analysis.

The Krawtchouk polynomials are important in the study of the Hamming scheme of classical coding theory [17], [21], [25], [29], [33], [34]. Lloyd's theorem [23] states that if a perfect code exists in the Hamming metric, then the Krawtchouk polynomial must have integral zeros [3], [5], [20]. Not surprisingly, these zeros have been the subject of extensive research [4], [7] [9], [10], [11], [16], [35].

The Krawtchouk polynomials also have applications in probability theory [8], queueing models [6], stochastic processes [31], quantum mechanics [2], [24] [39] and biology [13].

The asymptotic behavior of the Krawtchouk polynomials as $N \rightarrow \infty$ was studied by Sharapudinov for $x \approx Np$ and $n = O(N^{1/3})$ in [32]. He derived an approximation in terms of the Hermite polynomials (see also [36] for a similar formula). The general case with $x, n = O(N)$ was investigated by Ismail and Simeonov [12] and a uniform asymptotic expansion was derived by Li and Wong [22], both using the saddle point method.

The purpose of this paper is to take a different approach, based on the recurrence formula that the Krawtchouk polynomials satisfy and using singular perturbation techniques [26], [37] to analyze it. We scale $x = yN$, $n = zN$ and obtain asymptotic approximations to $K_n(x)$ for $(y, z) \in [0, 1] \times [0, 1]$. Our results agree and extend those obtained in [12], [22] and [30].

In Section 2 we review the basic properties of the Krawtchouk polynomials. In Sections 3-10 we obtain asymptotic expansions from the recurrence formula by using the WKB method. We must consider twelve relevant regions of the two-dimensional state space. In Section 11 we summarize our results and numerically compare our approximations with the exact formula.

2 The WKB approximation

To analyze the recurrence

$$(n+1)K_{n+1}(x) + pq(N-n+1)K_{n-1}(x) + [p(N-n) + nq - x]K_n(x) = 0 \quad (4)$$

subject to the boundary conditions

$$K_0(x) = 1 \quad (5)$$

$$K_{N+1}(x) = \binom{x}{N+1} \quad (6)$$

$$K_n(0) = \binom{N}{n} (-p)^n \quad (7)$$

$$K_n(N) = \binom{N}{n} q^n \quad (8)$$

for large N , we introduce the scaled variables y, z defined by

$$x = yN, \quad n = zN, \quad 0 < y, z < 1. \quad (9)$$

We define the function $F(y, z)$ and the small parameter ε by

$$\varepsilon = \frac{1}{N}, \quad K_n(x) = F(\varepsilon x, \varepsilon n) = F(y, z) \quad (10)$$

and observe that $K_{n\pm 1}(x) = F(y, z \pm \varepsilon)$.

Substituting (9)-(10) in (4) we get

$$(z + \varepsilon)F(y, z + \varepsilon) + pq(1 - z + \varepsilon)F(y, z - \varepsilon) + [p(1 - z) + zq - y]F(y, z) = 0. \quad (11)$$

To find $F(y, z)$ for ε small, we shall use the WKB method [28]. Thus, we consider solutions which have the asymptotic form

$$F(y, z) \sim \varepsilon^\nu \exp[\varepsilon^{-1}\psi(y, z)] L(y, z). \quad (12)$$

Using (12) in (11), with

$$\varepsilon^{-1}\psi(y, z \pm \varepsilon) = \varepsilon^{-1}\psi(y, z) \pm \psi_z(y, z) + \frac{1}{2}\psi_{zz}(y, z)\varepsilon + O(\varepsilon^2),$$

dividing by $\exp[\varepsilon^{-1}\psi(y, z)]$ and expanding in powers of ε we obtain the *eikonal equation*

$$zU^2 + [p - y + z(q - p)]U + pq(1 - z) = 0 \quad (13)$$

and the *transport equation*

$$[zU^2 - pq(1 - z)]L_z + \left\{ \frac{1}{2}[zU^2 + pq(1 - z)]\psi_{zz} + U^2 + pq \right\}L = 0 \quad (14)$$

with

$$U(y, z) = \exp[\psi_z(y, z)]. \quad (15)$$

2.1 The functions ψ and L

From (15) we have

$$\psi(y, z) = \int \ln [U(y, z)] dz = z \ln(U) - \int z \frac{U_z}{U} dz = z \ln(U) - \int z \frac{U_z}{U} \frac{dz}{dU} dU$$

and using (13) we get

$$\int z \frac{U_z}{U} \frac{dz}{dU} dU = \int \frac{(y-p)U - pq}{(U+q)(U-p)} dU = \int \left[\frac{1}{U} + \frac{y-1}{U} - \frac{y}{U+q} \right] dU.$$

Hence,

$$\psi(y, z) = \ln [U^{z-1} (U-p)^{1-y} (U+q)^y] + A(y) \quad (16)$$

where the function $A(y)$ is still unknown.

From (14) we have

$$L(y, z) = B(y) \exp \left[- \int \frac{\frac{1}{2} [zU^2 + pq(1-z)] \psi_{zz} + U^2 + pq}{zU^2 - pq(1-z)} dz \right]$$

and from (15) $\psi_{zz} = U_z/U$. After changing variables from z to U , we obtain

$$L(y, z) = B(y) \exp \left\{ \int \left[\frac{1}{2U} + \frac{1}{U+q} - \frac{(p-y)U + pq}{(p-y)U^2 + 2pqU + pq(q-y)} \right] dU \right\}.$$

Thus,

$$L(y, z) = B(y) (U+q) \sqrt{\frac{U}{(p-y)U^2 + 2pqU + pq(q-y)}} \quad (17)$$

where $B(y)$ is to be determined.

2.2 The function U

Rewriting (13) as

$$U^2 + \left(\frac{p-y}{z} + q - p \right) U + (U_0)^2 = 0$$

and solving for U we get

$$U^\pm(y, z) = -\frac{1}{2} \left(\frac{p-y}{z} + q - p \right) \pm \frac{1}{2} \sqrt{\left(\frac{p-y}{z} + q - p \right)^2 - 4(U_0)^2} \quad (18)$$

where

$$U_0(z) = \sqrt{\frac{pq(1-z)}{z}}. \quad (19)$$

The discriminant in (18) vanishes if $\frac{p-y}{z} + q - p = \pm 2U_0$, which is equivalent to $y = Y^\pm$, with

$$Y^\pm(z) = p + (q - p)z \pm 2zU_0. \quad (20)$$

Rewriting the equation $\left(\frac{p-y}{z} + q - p\right)^2 - 4(U_0)^2 = 0$ as

$$\left(y - \frac{1}{2}\right)^2 + \left(z - \frac{1}{2}\right)^2 + 2(p - q)\left(y - \frac{1}{2}\right)\left(z - \frac{1}{2}\right) - pq = 0 \quad (21)$$

we can see that

$$\{(y, z) \in [0, 1] \times [0, 1] : y = Y^\pm(z)\} = \mathbb{E} \quad (22)$$

where $\mathbb{E}$ is an ellipse centered at $(\frac{1}{2}, \frac{1}{2})$ (see Figure 1). After rotation by $\pm \frac{\pi}{4}$ and translation to the origin, $\mathbb{E}$ reduces to one of the canonical forms

$$\begin{aligned} \frac{\bar{y}^2}{q/2} + \frac{\bar{z}^2}{p/2} &= 1, & p < q \\ \bar{y}^2 + \bar{z}^2 &= \frac{1}{4}, & p = \frac{1}{2} = q \\ \frac{\bar{y}^2}{p/2} + \frac{\bar{z}^2}{q/2} &= 1, & p > q. \end{aligned}$$

The ellipse $\mathbb{E}$ is contained in the square $[0, 1] \times [0, 1]$ and intersects the y and z axis at the points $(0, p)$, $(1, q)$, $(p, 0)$ and $(q, 1)$. Its left side between the points $(q, 1)$ and $(p, 0)$ coincides with the curve $Y^-(z)$ and its right side with the curve $Y^+(z)$.

For points (y, z) located outside $\mathbb{E}$, $U^\pm(y, z)$ are real and for points (y, z) inside $\mathbb{E}$ $U^\pm(y, z)$ are complex conjugates. When $y = Y^\pm$ the two values $U^\pm$ coalesce and we have

$$U^+(Y^+, z) = U_0 = U^-(Y^+, z), \quad U^+(Y^-, z) = -U_0 = U^-(Y^-, z). \quad (23)$$

Writing the function $L(y, z)$ in terms of U_0 we have from (17)

$$L(y, z) = B(y) \sqrt{\frac{(U - p)(U + q)}{z(U^2 - U_0^2)}}. \quad (24)$$

In the rest of this paper we shall use the following notation

$$\psi^\pm(y, z) = \ln \left[(U^\pm)^{z-1} (U^\pm - p)^{1-y} (U^\pm + q)^y \right] \quad (25)$$

and

$$L^\pm(y, z) = \sqrt{\frac{(U^\pm - p)(U^\pm + q)}{z[(U^\pm)^2 - U_0^2]}}. \quad (26)$$

Hence, we write

$$\begin{aligned} K_n(x) &\sim \varepsilon^\nu B^-(y) \exp \left[\varepsilon^{-1} \psi^-(y, z) + \varepsilon^{-1} A^-(y) \right] L^-(y, z) \\ &\quad + \varepsilon^\nu B^+(y) \exp \left[\varepsilon^{-1} \psi^+(y, z) + \varepsilon^{-1} A^+(y) \right] L^+(y, z) \end{aligned} \quad (27)$$

where $B^\pm(y)$ and $A^\pm(y)$ are functions to be determined. From (26) we see that $L^\pm(y, z)$ are singular when $U^\pm = U_0$, i.e., for $y = Y^\pm$, and also for $z = 0$. Therefore, we need to find asymptotic solutions valid in those regions.




Figure 1: A sketch of the ellipse $\mathbb{E}$ and the curves $Y^\pm(z)$.

3 The boundary $n = 0$ (Region I)

For $n = O(1)$ we have from (1) that $K_n(x) = O(N^n)$ as $N \rightarrow \infty$. Thus, we introduce the function $R_n^{(1)}(y)$ and consider solutions of (4) which have the asymptotic form

$$K_n(x) = N^n R_n^{(1)}\left(\frac{x}{N}\right). \quad (28)$$

Using (28) in (4) and expanding in powers of N gives, to leading order

$$(n+1)R_{n+1}^{(1)} + (p-y)R_n^{(1)} = 0.$$

Solving the recursion above with the initial condition (5) we get

$$R_n^{(1)}(y) = \frac{(y-p)^n}{n!}$$

and hence

$$K_n(x) \sim K_n^{(1)}(y) = N^n \frac{(y-p)^n}{n!}, \quad n = O(1). \quad (29)$$

3.1 The corner layer at $(p, 0)$ (Region II)

We shall now find an asymptotic solution in the neighborhood of the point $(p, 0)$. We introduce the stretched variable η and the function $R_n^{(2)}(\eta)$ defined by

$$\begin{aligned} y &= p + \eta\sqrt{2pq\varepsilon} \quad \eta = O(1), \\ K_n(x) &= \frac{\varepsilon^{-n/2}}{n!} \left(\frac{pq}{2}\right)^{n/2} R_n^{(2)}\left(\frac{\varepsilon x - p}{\sqrt{2pq\varepsilon}}\right). \end{aligned} \quad (30)$$

Using (30) in (4) yields, to leading order, the equation

$$R_{n+1}^{(2)} - 2\eta R_n^{(2)} + 2nR_{n-1}^{(2)} = 0$$

which we recognize as the recurrence relation for the Hermite polynomials. Thus,

$$K_n(x) \sim K_n^{(2)}(\eta) = \frac{\varepsilon^{-n/2}}{n!} \left(\frac{pq}{2}\right)^{n/2} H_n(\eta) \quad (31)$$

for $n = O(1)$ and $y - p = O(\varepsilon^{1/2})$, where $H_n(\eta)$ is the Hermite polynomial of degree n .

4 The lower corners (Regions III and IV)

Setting $n = z/\varepsilon$ in (29) and letting $\varepsilon \rightarrow 0$ we obtain

$$K_n^{(1)}(y) \sim \varepsilon^{1/2} \frac{1}{\sqrt{2\pi z}} \exp\left\{\varepsilon^{-1}[1 - \ln(z) + \ln(y-p)]z\right\} \quad (32)$$

where we have used Stirling's formula [1]

$$\Gamma(x) \sim \sqrt{\frac{2\pi}{x}} x^x e^{-x}, \quad x \rightarrow \infty. \quad (33)$$

From (25)-(26) we have as $z \rightarrow 0$

$$\psi^-(y, z) \sim \begin{cases} [1 - \ln(z) + \ln(y - p)] z, & y < p \\ (1 - y) \ln\left(\frac{1-y}{q}\right) + y \ln\left(\frac{y}{p}\right) + \ln\left(\frac{pq}{y-p}\right) z, & y > p \end{cases} \quad (34)$$

$$L^-(y, z) = \begin{cases} z^{-1/2} + O(z), & y < p \\ \frac{\sqrt{(y-1)y}}{y-p} + O(z), & y > p \end{cases} \quad (35)$$

$$\psi^+(y, z) \sim \begin{cases} (1 - y) \ln\left(\frac{1-y}{q}\right) + y \ln\left(\frac{y}{p}\right) + \ln\left(\frac{pq}{y-p}\right) z, & y < p \\ [1 - \ln(z) + \ln(y - p)] z, & y > p \end{cases} \quad (36)$$

and

$$L^+(y, z) = \begin{cases} \frac{\sqrt{(y-1)y}}{p-y} + O(z), & y < p \\ z^{-1/2} + O(z), & y > p \end{cases}. \quad (37)$$

Matching (34)-(37) and (32) we conclude that

$$K_n(x) \sim \begin{cases} K^-(y, z), & 0 < y < Y^-(z) \\ K^+(y, z), & Y^+(z) < y < 1 \end{cases} \quad (38)$$

with

$$K^-(y, z) = \varepsilon^{1/2} \frac{1}{\sqrt{2\pi}} \exp[\varepsilon^{-1} \psi^-(y, z)] L^-(y, z) \quad (39)$$

$$K^+(y, z) = \varepsilon^{1/2} \frac{1}{\sqrt{2\pi}} \exp[\varepsilon^{-1} \psi^+(y, z)] L^+(y, z). \quad (40)$$

Remark 2 *In the remainder of the paper, we will find asymptotic formulas only in the region $0 \leq y \leq Y^+(z)$, $0 \leq z \leq 1$. The corresponding results for $Y^+(z) \leq y \leq 1$ can be obtained by using the symmetry formula (3) and noting that under the transformations $y \rightarrow 1 - y$, $p \leftrightarrow q$, $K_n \rightarrow (-1)^n K_n$, we obtain*

$$\begin{aligned} U^- &\rightarrow U^+ \\ \psi^- &\rightarrow \psi^+ + z\pi i, \quad L^- \rightarrow L^+ \\ K^- &\rightarrow K^+. \end{aligned}$$

5 The boundary $x = 0$

We shall now consider the case $x = O(1)$.

Lemma 3 *Let $x = m$, with m an integer, $m \ll N$.*

1.

$$K_n(m) = (-p)^n \binom{N}{n} \left(1 - \frac{n}{p} N^{-1}\right)^m, \quad m = 0, 1. \quad (41)$$

2. *If $n = O(N)$, then*

$$K_n(m) \sim (-p)^n \binom{N}{n} \left(1 - \frac{n}{p} N^{-1}\right)^m, \quad N \rightarrow \infty, \quad m \geq 2. \quad (42)$$

Proof. From (1) we have for $x = m$ integer

$$\begin{aligned} K_n(m) &= \sum_{k=0}^m \binom{m}{k} \binom{N-m}{n-k} q^k (-p)^{n-k} \\ &= (-p)^n \binom{N}{n} \sum_{k=0}^m \binom{m}{k} \frac{\binom{N-m}{n-k}}{\binom{N}{n}} \left(-\frac{q}{p}\right)^k \end{aligned} \quad (43)$$

and (41) follows for $m = 0, 1$.

Setting $n = zN$ and using (33) we get

$$\frac{\binom{N-m}{Nz-k}}{\binom{N}{Nz}} \sim (1-z)^m \left(\frac{z}{1-z}\right)^k, \quad N \rightarrow \infty.$$

Using the above in (43) we have

$$\begin{aligned} K_n(m) &\sim (-p)^n \binom{N}{n} (1-z)^m \sum_{k=0}^m \binom{m}{k} \left(-\frac{z}{1-z} \frac{q}{p}\right)^k \\ &= (-p)^n \binom{N}{n} (1-z)^m \left[\frac{p-z}{p(1-z)}\right]^m \end{aligned}$$

and (42) follows. ■

Using (33) we have, as $N \rightarrow \infty$

$$(-p)^n \binom{N}{n} \sim \frac{\varepsilon^{1/2}}{\sqrt{2\pi} \sqrt{z(1-z)}} \exp[\varepsilon^{-1} \phi_0(z)] \quad (44)$$

where

$$\phi_0(z) = (z-1) \ln(1-z) - z \ln(z) + z \ln(-p). \quad (45)$$

From (39) we get, as $y \rightarrow 0$

$$K^-(y, z) \sim \frac{\varepsilon^{1/2}}{\sqrt{2\pi}\sqrt{z(1-z)}} \exp \left[\frac{\phi_0(z) + \ln \left(\frac{p-z}{p} \right) y}{\varepsilon} \right], \quad 0 < z < p. \quad (46)$$

Hence, as $y \rightarrow 0$, $K^-(y, z)$ satisfies the boundary condition (7) for $0 < z < p$.

5.1 The boundary layer at $x = 0$, $p < z < 1$ (Region V)

Taking (42) into account, we define the function $R_n^{(5)}(x)$ by

$$K_n(x) = \binom{N}{n} (-p)^n R_n^{(5)}(x). \quad (47)$$

Using (47) in (4) yields

$$p(n-N)R_{n+1}^{(5)} + [pN + (q-p)n - x] R_n^{(5)} - nqR_{n-1}^{(5)} = 0. \quad (48)$$

Writing (48) in terms of z, ε and the function $G^{(5)}(x, z)$ defined by

$$R_n^{(5)}(x) = G^{(5)}(x, \varepsilon n) \quad (49)$$

we have

$$p(1-z)G^{(5)}(x, z+\varepsilon) + [z(p-q) + x\varepsilon - p] G^{(5)}(x, z) + zqG^{(5)}(x, z-\varepsilon) = 0. \quad (50)$$

Using the WKB ansatz

$$G^{(5)}(x, z) \sim \varepsilon^\tau \exp \left[\frac{1}{\varepsilon} \phi(x, z) \right] W(x, z) \quad (51)$$

in (50), we obtain the equations

$$[\exp(\phi_z) - 1] [p(z-1) \exp(\phi_z) + zq] W = 0 \quad (52)$$

and

$$\begin{aligned} & [zq + p(z-1) \exp(2\phi_z)] W_z \\ & + \left\{ \frac{1}{2} [p(z-1) \exp(2\phi_z) - zq] \phi_{zz} - x \exp(\phi_z) \right\} W = 0. \end{aligned} \quad (53)$$

Solving (52)-(53) we get

$$\begin{aligned} G^{(5)}(x, z) & \sim \varepsilon^{\tau_1} A^{(5)}(x) \exp [\varepsilon^{-1} \varphi_1(x)] (z-p)^x \\ & + \varepsilon^{\tau_2} B^{(5)}(x) \exp [\varepsilon^{-1} \phi_2(x, z)] (z-p)^{-x-1} \sqrt{z(1-z)} \end{aligned} \quad (54)$$

where

$$\phi_2(x, z) = (1 - z) \ln(1 - z) + z \ln(z) + (z - 1) \ln(q) - z \ln(p) + \varphi_2(x) \quad (55)$$

and the coefficients τ_1, τ_2 and the functions $A^{(5)}(x), B^{(5)}(x), \varphi_1(x), \varphi_2(x)$ are to be determined.

Using (44) we obtain

$$\begin{aligned} K_n(x) \sim K^{(5)}(x, z) &= \frac{\varepsilon^{1/2+\tau_1}}{\sqrt{2\pi}\sqrt{z(1-z)}} \exp\left[\frac{\phi_0(z) + \varphi_1(x)}{\varepsilon}\right] A(x) (z - p)^x \\ &+ \frac{\varepsilon^{1/2+\tau_2}}{\sqrt{2\pi}} \exp\left[\frac{(z - 1) \ln(q) + \pi i z + \varphi_2(x)}{\varepsilon}\right] B(x) (z - p)^{-x-1}. \end{aligned} \quad (56)$$

5.2 The corner layer at $(0, p)$ (Region VI)

For $x \approx 0$ and $n \approx Np$, we scale n as

$$n = Np - u\sqrt{pqN}, \quad u = O(1) \quad (57)$$

and introduce the function $G^{(6)}(x, u)$ defined by

$$K_n(x) = \binom{N}{n} (-p)^n G^{(6)}\left(x, \frac{Np - n}{\sqrt{pqN}}\right). \quad (58)$$

Using (57)-(58) in (48) we get, to leading order

$$\frac{\partial^2 G^{(6)}}{\partial u^2} - \frac{\partial G^{(6)}}{\partial u} + x G^{(6)} = 0. \quad (59)$$

Solving (59) we obtain

$$G^{(6)}(x, u) = \exp\left(\frac{u^2}{4}\right) [A^{(6)}(x) D_x(u) + B^{(6)}(x) D_x(-u)] \quad (60)$$

where $D_x(u)$ is the parabolic cylinder function and $A^{(6)}(x), B^{(6)}(x)$ are functions to be determined. Since

$$u = \frac{Np - n}{\sqrt{pqN}} = \frac{p - z}{\sqrt{pq\varepsilon}}, \quad (61)$$

we note that the limit $u \rightarrow \infty$ corresponds to the matching between regions VI and III, while the limit $u \rightarrow -\infty$ corresponds to the matching between regions VI and V. As $z \rightarrow p$ we have

$$\begin{aligned} (-p)^n \binom{N}{n} &\sim \frac{\sqrt{\varepsilon}}{\sqrt{2\pi pq}} \exp\left(-\frac{u^2}{2}\right) \\ &\times \exp\left\{\frac{\pi i p - q \ln(q)}{\varepsilon} - \frac{u\sqrt{pq}[\pi i + \ln(q)]}{\sqrt{\varepsilon}}\right\}. \end{aligned} \quad (62)$$

Thus,

$$K_n(x) \sim K^{(6)}(x, u) = \frac{\sqrt{\varepsilon}}{\sqrt{2\pi pq}} [A^{(6)}(x)D_x(u) + B^{(6)}(x)D_x(-u)] \quad (63)$$

$$\times \exp \left\{ \frac{\pi i p - q \ln(q)}{\varepsilon} - \frac{u\sqrt{pq} [\pi i + \ln(q)]}{\sqrt{\varepsilon}} - \frac{u^2}{4} \right\}.$$

Using (61) in (46) yields

$$K^-(y, z) \sim \frac{\sqrt{\varepsilon}}{\sqrt{2\pi pq}} \left[\sqrt{\frac{q\varepsilon}{p}} u \right]^x \quad (64)$$

$$\times \exp \left[\frac{\pi i p - q \ln(q)}{\varepsilon} - \frac{u\sqrt{pq} [\pi i + \ln(q)]}{\sqrt{\varepsilon}} - \frac{u^2}{2} \right].$$

Using the well known asymptotic approximation [38]

$$D_x(u) \sim \exp \left(-\frac{u^2}{4} \right) u^x, \quad u \rightarrow \infty \quad (65)$$

in (63) and comparing with (64) we conclude that

$$A^{(6)}(x) = \left[\sqrt{\frac{q\varepsilon}{p}} \right]^x$$

and $B^{(6)}(x) = 0$. Therefore,

$$K^{(6)}(x, u) = \frac{\varepsilon^{1/2}}{\sqrt{2\pi pq}} \left[\sqrt{\frac{q\varepsilon}{p}} \right]^x D_x(u) \quad (66)$$

$$\times \exp \left[\frac{\pi i p - q \ln(q)}{\varepsilon} - \frac{u\sqrt{pq} [\pi i + \ln(q)]}{\sqrt{\varepsilon}} - \frac{u^2}{4} \right]$$

for $x = O(1)$, $z - p = O(\varepsilon^{1/2})$.

Using the formula [38]

$$D_x(-u) \sim \exp \left(-\frac{u^2}{4} \right) u^x \cos(\pi x) - \sqrt{\frac{2}{\pi}} x \Gamma(x) \sin(\pi x) u^{-x-1} \exp \left(\frac{u^2}{4} \right), \quad u \rightarrow \infty \quad (67)$$

in (66) we have

$$K^{(6)}(x, -u) \sim \frac{\varepsilon^{1/2}}{\sqrt{2\pi pq}} \left[u \sqrt{\frac{q\varepsilon}{p}} \right]^x \cos(\pi x)$$

$$\times \exp \left[\frac{\pi i p - q \ln(q)}{\varepsilon} - \frac{u\sqrt{pq} [\pi i + \ln(q)]}{\sqrt{\varepsilon}} - \frac{u^2}{2} \right] \quad (68)$$

$$- \frac{1}{u} \frac{\varepsilon^{1/2}}{\pi \sqrt{pq}} \left[\frac{1}{u} \sqrt{\frac{q\varepsilon}{p}} \right]^x x \Gamma(x) \sin(\pi x)$$

$$\times \exp \left[\frac{\pi i p - q \ln(q)}{\varepsilon} - \frac{u\sqrt{pq} [\pi i + \ln(q)]}{\sqrt{\varepsilon}} \right], \quad u \rightarrow \infty.$$

Using (61) in (56) gives, as $z \downarrow p$

$$\begin{aligned}
K^{(5)}(x, z) &\sim \frac{\varepsilon^{1/2+\tau_1}}{\sqrt{2\pi pq}} A^{(5)}(x) (u\sqrt{pq\varepsilon})^x \\
&\times \exp \left[\frac{\pi i p - q \ln(q) + \varphi_1(x)}{\varepsilon} - \frac{u\sqrt{pq} [\pi i + \ln(q)]}{\sqrt{\varepsilon}} - \frac{u^2}{2} \right] \\
&+ \exp \left[\frac{\pi i p - q \ln(q) + \varphi_2(x)}{\varepsilon} - \frac{u\sqrt{pq} [\pi i + \ln(q)]}{\sqrt{\varepsilon}} \right] \\
&\times \frac{1}{u} \frac{\varepsilon^{\tau_2}}{\sqrt{2\pi pq}} B^{(5)}(x) (u\sqrt{pq\varepsilon})^{-x}.
\end{aligned} \tag{69}$$

Matching (68) and (69) yields $\tau_1 = 0$, $\varphi_1(x) = 0$, $\tau_2 = \frac{1}{2}$, $\varphi_2(x) = 0$ and

$$\begin{aligned}
A^{(5)}(x) &= p^{-x} \cos(\pi x) \\
B^{(5)}(x) &= -\sqrt{\frac{2}{\pi}} x \Gamma(x) \sin(\pi x) (q\varepsilon)^x.
\end{aligned}$$

Thus,

$$\begin{aligned}
K^{(5)}(x, z) &= \frac{\varepsilon^{1/2}}{\sqrt{2\pi} \sqrt{z(1-z)}} \cos(\pi x) \left(\frac{z-p}{p} \right)^x \exp \left[\frac{\phi_0(z)}{\varepsilon} \right] \\
&- \frac{\varepsilon}{\pi} \frac{x}{z-p} \Gamma(x) \sin(\pi x) \left(\frac{q\varepsilon}{z-p} \right)^x \exp \left[\frac{(z-1) \ln(q) + \pi i z}{\varepsilon} \right]
\end{aligned} \tag{70}$$

for $x = O(1)$ and $p < z < 1$.

From (70) we have, as $x \rightarrow \infty$

$$\begin{aligned}
K^{(5)}(x, z) &\sim \frac{\varepsilon^{1/2}}{\sqrt{2\pi} \sqrt{z(1-z)}} \cos(\pi x) \left(\frac{z-p}{p} \right)^x \exp \left[\frac{\phi_0(z)}{\varepsilon} \right] \\
&- \varepsilon \sqrt{\frac{2}{\pi}} \frac{\sqrt{x}}{z-p} \sin(\pi x) \left(\frac{q\varepsilon x}{z-p} \right)^x \exp \left[\frac{(z-1) \ln(q) + \pi i z}{\varepsilon} - x \right].
\end{aligned} \tag{71}$$

In terms of $y = x\varepsilon$, (71) reads

$$\begin{aligned}
K^{(5)}(x, z) &\sim \frac{\varepsilon^{1/2}}{\sqrt{2\pi} \sqrt{z(1-z)}} \cos \left(\frac{\pi y}{\varepsilon} \right) \exp \left[\frac{\phi_0(z) + y \ln \left(\frac{z-p}{p} \right)}{\varepsilon} \right] \\
&- \varepsilon^{1/2} \sqrt{\frac{2}{\pi}} \sin \left(\frac{\pi y}{\varepsilon} \right) \frac{\sqrt{y}}{z-p} \exp \left[\frac{(z-1) \ln(q) + \pi i z + y \ln \left(\frac{qy}{z-p} \right) - y}{\varepsilon} \right].
\end{aligned} \tag{72}$$

6 The left upper corner (Region VII)

We now consider the region $0 \ll y < Y^-(z)$, $p < z < 1$. From (25)-(26) we have for $y \rightarrow 0$, $z > p$

$$\psi^+(y, z) \sim \phi_0(z) + y \ln \left(\frac{z-p}{p} \right) - \pi i y, \quad (73)$$

$$\psi^-(y, z) \sim (z-1) \ln(q) + \pi i z + y \ln \left(\frac{qy}{z-p} \right) - y - \pi i y, \quad (74)$$

and

$$L^+(y, z) \sim \frac{1}{\sqrt{z(1-z)}}, \quad L^-(y, z) \sim \frac{\sqrt{y}}{z-p} i. \quad (75)$$

Using (73)-(75) in (27) and matching with (72), we conclude that

$$K_n(x) \sim K^{(7)}(y, z) = \exp \left(\frac{\pi i y}{\varepsilon} \right) \left[\cos \left(\frac{\pi y}{\varepsilon} \right) K^+(y, z) + 2i \sin \left(\frac{\pi y}{\varepsilon} \right) K^-(y, z) \right], \quad (76)$$

which can be written as

$$K^{(7)}(y, z) = \frac{1}{2} \left[\exp \left(\frac{2\pi i y}{\varepsilon} \right) + 1 \right] K^+(y, z) + \left[\exp \left(\frac{2\pi i y}{\varepsilon} \right) - 1 \right] K^-(y, z). \quad (77)$$

7 The transition layer $y = Y^-(z)$

As we noted before, the functions $L^\pm(y, z)$ are infinite on the curves $y = Y^\pm(z)$. Hence, we need to find transition layer solutions there.

7.1 The lower part $0 < z < p$ (Region VIII)

We introduce the stretched variable β , defined by

$$y = Y^-(z) - \beta \varepsilon^{2/3} \quad \beta = O(1), \quad 0 < z < p. \quad (78)$$

Using (78) in (25)-(26) and expanding in powers of ε , with $\beta > 0$, we get

$$\psi^-(y, z) \sim \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{2/3} - \frac{2}{3} \sqrt{\frac{U_0}{z}} \frac{1}{(U_0 + p)(U_0 - q)} \beta^{3/2} \varepsilon \quad (79)$$

where

$$\psi_0(z) = z\pi i + (z-1) \ln(U_0) + Y^-(z) \ln(U_0 - q) + [1 - Y^-(z)] \ln(U_0 + p) \quad (80)$$

and

$$L^-(y, z) \sim \frac{1}{\sqrt{2}} z^{-1/4} \sqrt{(U_0 + p)(U_0 - q)} (U_0)^{-3/4} \beta^{-1/4} \varepsilon^{-1/6}. \quad (81)$$

Hence,

$$K^-(y, z) \sim \varepsilon^{1/3} \frac{1}{2\sqrt{\pi}} z^{-1/4} \sqrt{(U_0 + p)(U_0 - q)} (U_0)^{-3/4} \beta^{-1/4} \quad (82)$$

$$\times \exp \left[\varepsilon^{-1} \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{-1/3} - \frac{2}{3} \sqrt{\frac{U_0}{z}} \frac{1}{(U_0 + p)(U_0 - q)} \beta^{3/2} \right]$$

for $y \uparrow Y^-(z)$. Thus, we introduce the function $G^{(8)}(\beta, z)$ and consider solutions of the form

$$K_n(x) = \varepsilon^{\nu_8} \exp \left[\psi_0(z) \varepsilon^{-1} + \ln \left(\frac{U_0 + p}{U_0 - q} \right) (Y^- - \varepsilon x) \varepsilon^{-1} \right] G^{(8)} \left(\frac{Y^- - \varepsilon x}{\varepsilon^{2/3}}, \varepsilon n \right). \quad (83)$$

Using (83) in (4) and expanding in powers of ε we obtain

$$G_{\beta\beta}^{(8)} = - \frac{2\beta}{[(Y^-)]^2 [(p - q)z + Y^- - p]} G^{(8)} = \beta \Theta^2 G^{(8)} \quad (84)$$

where

$$\Theta(z) = \sqrt{\frac{U_0}{z}} \frac{1}{(U_0 + p)(U_0 - q)}. \quad (85)$$

Solving (84) we get

$$G^{(8)}(\beta, z) = A^{(8)}(z) \text{Ai} [\Theta^{2/3} \beta] + B^{(8)}(z) \text{Bi} [\Theta^{2/3} \beta] \quad (86)$$

where $A^{(8)}(z), B^{(8)}(z)$ are functions to be determined and $\text{Ai}(\cdot), \text{Bi}(\cdot)$ are the Airy functions. Using the formulas [1]

$$\text{Ai}(x) \sim \frac{1}{2\sqrt{\pi}} x^{-1/4} \exp \left(-\frac{2}{3} x^{3/2} \right), \quad x \rightarrow \infty \quad (87)$$

$$\text{Bi}(x) \sim \frac{1}{\sqrt{\pi}} x^{-1/4} \exp \left(\frac{2}{3} x^{3/2} \right), \quad x \rightarrow \infty$$

in (86) and matching with (82) we conclude that $\nu_8 = 1/3$, $B^{(8)}(z) = 0$ and

$$A^{(8)}(z) = \left[\frac{(U_0 + p)(U_0 - q)}{z(U_0)^2} \right]^{1/3} = \frac{\Theta^{-1/3}}{\sqrt{zU_0}}. \quad (88)$$

Therefore,

$$K^{(8)}(\beta, z) = \varepsilon^{1/3} \exp \left[\varepsilon^{-1} \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{-1/3} \right] \text{Ai} [\Theta^{2/3} \beta] \frac{\Theta^{-1/3}}{\sqrt{zU_0}} \quad (89)$$

for $y - Y^-(z) = O(\varepsilon^{2/3})$.

7.2 The upper part $p < z < 1$ (Region IX)

Using (78) in (25) we have, as $y \uparrow Y^-(z)$

$$\psi^-(y, z) \sim \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{2/3} + \frac{2}{3} \vartheta \beta^{3/2} \varepsilon, \quad (90)$$

$$\psi^+(y, z) \sim \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{2/3} - \frac{2}{3} \vartheta \beta^{3/2} \varepsilon \quad (91)$$

and from (26) we have

$$L^-(y, z) \sim \frac{1}{\sqrt{2}} \frac{1}{\sqrt{z U_0 \vartheta}} i \beta^{-1/4} \varepsilon^{-1/6}, \quad (92)$$

$$L^+(y, z) \sim \frac{1}{\sqrt{2}} \frac{1}{\sqrt{z U_0 \vartheta}} \beta^{-1/4} \varepsilon^{-1/6} \quad (93)$$

where

$$\vartheta(z) = -\Theta(z) = \sqrt{\frac{U_0}{z}} \frac{1}{(U_0 + p)(q - U_0)}. \quad (94)$$

Using (90)-(93) in (77) we have

$$\begin{aligned} K^{(7)}(y, z) &\sim \frac{1}{2\sqrt{\pi}} \frac{1}{\sqrt{z U_0 \vartheta}} \beta^{-1/4} \varepsilon^{1/3} \exp \left[\psi_0(z) \varepsilon^{-1} + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{-1/3} \right] \\ &\times \left[\frac{1}{2} \lambda^+(\beta, z) \exp \left(-\frac{2}{3} \vartheta \beta^{3/2} \right) + i \lambda^-(\beta, z) \exp \left(\frac{2}{3} \vartheta \beta^{3/2} \right) \right] \end{aligned} \quad (95)$$

as $y \uparrow Y^-(z)$, where

$$\lambda^\pm(\beta, z) = \exp \left\{ \frac{2\pi i [Y^-(z) - \beta \varepsilon^{2/3}]}{\varepsilon} \right\} \pm 1. \quad (96)$$

We consider the ansatz

$$\begin{aligned} K_n(x) &\sim K^{(9)}(\beta, z) = \varepsilon^{\nu_9} \exp \left[\varepsilon^{-1} \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{-1/3} \right] \\ &\times [\lambda^+(\beta, z) A^{(9)}(z) \text{Ai}(\vartheta^{2/3} \beta) + \lambda^-(\beta, z) B^{(9)}(z) \text{Bi}(\vartheta^{2/3} \beta)] \end{aligned} \quad (97)$$

for $y \approx Y^-(z)$, $p < z < 1$ and unknown functions $A^{(9)}(z), B^{(9)}(z)$. Using (87) in (97) we have, as $\beta \rightarrow \infty$

$$\begin{aligned} K^{(9)}(\beta, z) &\sim \varepsilon^{\nu_9} \exp \left[\varepsilon^{-1} \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{-1/3} \right] \\ &\times \left[\lambda^+(\beta, z) A^{(9)}(z) \frac{1}{2\sqrt{\pi}} \vartheta^{-1/6} \beta^{-1/4} \exp \left(-\frac{2}{3} \vartheta \beta^{3/2} \right) + \right. \\ &\left. \lambda^-(\beta, z) B^{(9)}(z) \frac{1}{\sqrt{\pi}} \vartheta^{-1/6} \beta^{-1/4} \exp \left(\frac{2}{3} \vartheta \beta^{3/2} \right) \right]. \end{aligned} \quad (98)$$

Matching (98) and (95) we get

$$\nu_9 = \frac{1}{3}, \quad A^{(9)}(z) = \frac{1}{2} \frac{\vartheta^{-1/3}}{\sqrt{zU_0}}, \quad B^{(9)}(z) = \frac{1}{2} i \frac{\vartheta^{-1/3}}{\sqrt{zU_0}}.$$

Thus,

$$\begin{aligned} K^{(9)}(\beta, z) &= \varepsilon^{1/3} \exp \left[\varepsilon^{-1} \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{-1/3} \right] \\ &\times \frac{1}{2} \frac{\vartheta^{-1/3}}{\sqrt{zU_0}} \left[\lambda^+(\beta, z) \text{Ai}(\vartheta^{2/3} \beta) + i \lambda^-(\beta, z) \text{Bi}(\vartheta^{2/3} \beta) \right]. \end{aligned} \quad (99)$$

8 The interior of E (Region X)

We shall now find an asymptotic solution for $Y^-(z) < y < Y^+(z)$. We set $\beta = -\tilde{\beta}$, $\tilde{\beta} > 0$ in (79)-(81) and obtain, for $0 < z < p$

$$\psi^-(y, z) \sim \psi_0(z) - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{2/3} + \frac{2}{3} \Theta i \tilde{\beta}^{3/2} \varepsilon \quad (100)$$

and

$$L^-(y, z) \sim \frac{1}{\sqrt{2}} \frac{1}{\sqrt{zU_0\Theta}} e^{-\frac{1}{4}\pi i \tilde{\beta}^{-1/4} \varepsilon^{-1/6}}. \quad (101)$$

Similarly, from (25)-(26) we obtain, for $0 < z < p$

$$\psi^+(y, z) \sim \psi_0(z) - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{2/3} - \frac{2}{3} \Theta i \tilde{\beta}^{3/2} \varepsilon \quad (102)$$

and

$$L^+(y, z) \sim \frac{1}{\sqrt{2}} \frac{1}{\sqrt{zU_0\Theta}} e^{\frac{1}{4}\pi i \tilde{\beta}^{-1/4} \varepsilon^{-1/6}}. \quad (103)$$

Introducing the function $K^{(10)}(y, z)$ defined by

$$\begin{aligned} K^{(10)}(y, z) &= \varepsilon^{\nu_{10}} A^{(10)}(y) \exp \left[\varepsilon^{-1} \psi^-(y, z) \right] L^-(y, z) \\ &+ \varepsilon^{\nu_{10}} B^{(10)}(y) \exp \left[\varepsilon^{-1} \psi^+(y, z) \right] L^+(y, z) \end{aligned} \quad (104)$$

with $A^{(10)}(y), B^{(10)}(y)$ to be determined, we have

$$\begin{aligned} K^{(10)}(y, z) &\sim \varepsilon^{\nu_{10}-1/6} \frac{1}{\sqrt{2}} \frac{1}{\sqrt{zU_0\Theta}} \tilde{\beta}^{-1/4} \exp \left[\psi_0^-(z) \varepsilon^{-1} - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{-1/3} \right] \\ &\times \left[A^{(10)}(y) \exp \left(\frac{2}{3} \Theta i \tilde{\beta}^{3/2} - \frac{1}{4} \pi i \right) + B^{(10)}(y) \exp \left(-\frac{2}{3} \Theta i \tilde{\beta}^{3/2} + \frac{1}{4} \pi i \right) \right]. \end{aligned} \quad (105)$$

From (89) we have

$$K^{(9)}(\beta, z) \sim \varepsilon^{1/3} \frac{1}{\sqrt{\pi}} \frac{1}{\sqrt{zU_0\Theta}} \tilde{\beta}^{-1/4} \exp \left[\varepsilon^{-1} \psi_0^-(z) - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{-1/3} \right] \times \sin \left(\frac{2}{3} \Theta \tilde{\beta}^{3/2} + \frac{\pi}{4} \right) \quad (106)$$

where we have used the asymptotic formula [1]

$$\text{Ai}(-x) \sim \frac{x^{-1/4}}{\sqrt{\pi}} \sin \left(\frac{2}{3} x^{3/2} + \frac{\pi}{4} \right), \quad x \rightarrow \infty. \quad (107)$$

Matching (105) and (106) we get $\nu_{10} = \frac{1}{2}$ and

$$\begin{aligned} & \frac{1}{\sqrt{2}} \left[A^{(10)}(y) \exp \left(\frac{2}{3} \Theta i \tilde{\beta}^{3/2} - \frac{1}{4} \pi i \right) + B^{(10)}(y) \exp \left(-\frac{2}{3} \Theta i \tilde{\beta}^{3/2} + \frac{1}{4} \pi i \right) \right] \\ &= \frac{1}{\sqrt{\pi}} \sin \left(\frac{2}{3} \Theta \tilde{\beta}^{3/2} + \frac{\pi}{4} \right) = \frac{1}{2\sqrt{\pi}} \left[\exp \left(\frac{2}{3} \Theta i \tilde{\beta}^{3/2} - \frac{1}{4} \pi i \right) + \exp \left(-\frac{2}{3} \Theta i \tilde{\beta}^{3/2} + \frac{1}{4} \pi i \right) \right] \end{aligned}$$

from which we conclude that

$$A^{(10)}(y) = \frac{1}{\sqrt{2\pi}} = B^{(10)}(y)$$

and therefore

$$K_n(x) \sim K^{(10)}(y, z) = K^+(y, z) + K^-(y, z) \quad (108)$$

for $Y^-(z) < y < Y^+(z)$.

8.1 Matching the interior of E and the upper part of the transition layer Y^-

We shall now verify the matching between (108) and (99). For $p < z < 1$ we have

$$\psi^-(y, z) \sim (Y^- + \tilde{\beta} \varepsilon^{2/3} - z) 2\pi i + \psi_0(z) - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{2/3} - \frac{2}{3} \vartheta i \tilde{\beta}^{3/2} \varepsilon, \quad (109)$$

$$L^-(y, z) \sim \frac{1}{\sqrt{2}} \frac{1}{\sqrt{zU_0\vartheta}} e^{\frac{1}{4}\pi i} \tilde{\beta}^{-1/4} \varepsilon^{-1/6}, \quad (110)$$

$$\psi^+(y, z) \sim \psi_0(z) - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{2/3} + \frac{2}{3} \vartheta i \tilde{\beta}^{3/2} \varepsilon \quad (111)$$

and

$$L^+(y, z) \sim \frac{1}{\sqrt{2}} \frac{1}{\sqrt{zU_0\vartheta}} e^{-\frac{1}{4}\pi i} \tilde{\beta}^{-1/4} \varepsilon^{-1/6}. \quad (112)$$

Therefore,

$$K^{(10)}(y, z) \sim \frac{\varepsilon^{1/3}}{2\sqrt{\pi}} \frac{\tilde{\beta}^{-1/4}}{\sqrt{zU_0\vartheta}} \exp \left[\varepsilon^{-1} \psi_0(z) - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{-1/3} \right] \quad (113)$$

$$\times \left[\frac{\lambda^+(-\tilde{\beta}, z) + \lambda^-(-\tilde{\beta}, z)}{2} \exp \left(-\frac{2}{3} \vartheta i \tilde{\beta}^{3/2} + \frac{1}{4} \pi i \right) + \exp \left(\frac{2}{3} \vartheta i \tilde{\beta}^{3/2} - \frac{1}{4} \pi i \right) \right].$$

Using (107) and [1]

$$\text{Bi}(-x) \sim \frac{x^{-1/4}}{\sqrt{\pi}} \cos \left(\frac{2}{3} x^{3/2} + \frac{\pi}{4} \right), \quad x \rightarrow \infty$$

in (99) we have

$$K^{(9)}(\beta, z) \sim \varepsilon^{1/3} \exp \left[\varepsilon^{-1} \psi_0(z) - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{-1/3} \right]$$

$$\times \frac{1}{2\sqrt{\pi}} \frac{\tilde{\beta}^{-1/4}}{\sqrt{zU_0\vartheta}} \left[\lambda^+(-\tilde{\beta}, z) \sin \left(\frac{2}{3} \vartheta \tilde{\beta}^{3/2} + \frac{\pi}{4} \right) + i \lambda^-(-\tilde{\beta}, z) \cos \left(\frac{2}{3} \vartheta \tilde{\beta}^{3/2} + \frac{\pi}{4} \right) \right]$$

or

$$K^{(9)}(\beta, z) \sim \varepsilon^{1/3} \exp \left[\varepsilon^{-1} \psi_0(z) - \ln \left(\frac{U_0 + p}{U_0 - q} \right) \tilde{\beta} \varepsilon^{-1/3} \right] \frac{1}{2\sqrt{\pi}} \frac{\tilde{\beta}^{-1/4}}{\sqrt{zU_0\vartheta}} \quad (114)$$

$$\times \left[\frac{\lambda^+ + \lambda^-}{2} \exp \left(-\frac{2}{3} \vartheta i \tilde{\beta}^{3/2} + \frac{1}{4} \pi i \right) + \frac{\lambda^+ - \lambda^-}{2} \exp \left(\frac{2}{3} \vartheta i \tilde{\beta}^{3/2} - \frac{1}{4} \pi i \right) \right].$$

Since from (96) we have

$$\frac{\lambda^+ - \lambda^-}{2} = 1$$

we see that (114) agrees with (113).

8.2 Matching the interior of E and the corner layer at $(p, 0)$

Using (33) and the asymptotic formula [19]

$$H_n(x) \sim \sqrt{2}^{n+1} n^{n/2} \exp \left(\frac{x^2}{2} - \frac{n}{2} \right) \cos \left(\sqrt{2n}x - \frac{n\pi}{2} \right), \quad n \rightarrow \infty$$

in (31) we have, as $n \rightarrow \infty$

$$K_n^{(2)}(\eta) \sim \frac{1}{\sqrt{n\pi}} \exp \left\{ \frac{\eta^2}{2} + \frac{n}{2} \left[1 + \ln \left(\frac{pq}{\varepsilon n} \right) \right] \right\} \cos \left(\sqrt{2n}\eta - \frac{n\pi}{2} \right). \quad (115)$$

Using (30) in (25), with $z = n\varepsilon$ we get

$$\frac{1}{\varepsilon}\psi^\pm \sim \frac{n}{2} \left[1 + \ln \left(\frac{pq}{\varepsilon n} \right) \pm \pi i \right] \mp \sqrt{2n\eta}i + \frac{\eta^2}{2}. \quad (116)$$

Similarly from (26) we find

$$L^+ \sim \frac{1}{\sqrt{2n}}, \quad L^- \sim \frac{1}{\sqrt{2n}}. \quad (117)$$

Using (116)-(117) in (108) we obtain

$$K^{(10)} \sim \frac{1}{\sqrt{\pi n}} \exp \left\{ \frac{\eta^2}{2} + \frac{n}{2} \left[1 + \ln \left(\frac{pq}{\varepsilon n} \right) \right] \right\} \cos \left(\sqrt{2n\eta} - \frac{n\pi}{2} \right)$$

which is in agreement with (115).

9 The boundary layer at $z = 1$ (Region XI)

We now consider solutions of (4) with $n \approx N$. We introduce the variable j and the function $\Upsilon_j^{(11)}(y)$ defined by

$$n = N - j, \quad j \in \mathbb{Z}, \quad j \geq -1, \quad (118)$$

$$K_n(x) = \binom{N}{N-n} (-p)^n \Upsilon_{N-n}^{(11)} \left(\frac{x}{N} \right). \quad (119)$$

Using (119) in (4) we have, to leading order,

$$q\Upsilon_{j+1}^{(11)} - (q-y)\Upsilon_j^{(11)} = 0$$

which can be solved to obtain

$$\Upsilon_j^{(11)}(y) = A^{(11)}(y) \left(1 - \frac{y}{q} \right)^j \quad (120)$$

where $A^{(11)}(y)$ is a function to be determined. Thus, we have

$$K_n(x) \sim A^{(11)}(y) \binom{N}{j} (-p)^{N-j} \left(1 - \frac{y}{q} \right)^j. \quad (121)$$

We now introduce the function $R_j^{(11)}(y)$ defined by

$$K_n(x) = \binom{x}{n} R_{N-n}^{(11)} \left(\frac{x}{N} \right) \quad (122)$$

and note that from (??) we have

$$R_{-1}^{(11)}(y) = 1. \quad (123)$$

Using (118) in (4) we obtain, to leading order,

$$(q - y)R_{j+1}^{(11)} - (1 - y)R_j^{(11)} = 0$$

which together with (123) implies

$$R_j^{(11)}(y) = \left(\frac{1 - y}{q - y} \right)^{j+1}, \quad y \neq q. \quad (124)$$

Therefore,

$$K_n(x) \sim \binom{Ny}{n} \left(\frac{1 - y}{q - y} \right)^{j+1}, \quad y \not\approx q. \quad (125)$$

From (122) and (125) we get, for $y \not\approx q$,

$$K_n(x) \sim K_j^{(11)}(y) = A^{(11)}(y) \binom{N}{j} (-p)^{N-j} \left(1 - \frac{y}{q} \right)^j + \binom{Ny}{N-j} \left(\frac{1 - y}{q - y} \right)^{j+1}. \quad (126)$$

9.1 Matching with the upper corners

We shall now determine the function $A^{(11)}(y)$ by matching (126) with (76). As $N \rightarrow \infty$ we get

$$\begin{aligned} K_j^{(11)}(y) \sim A^{(11)}(y) \frac{1}{\sqrt{2\pi j}} \exp \left\{ \varepsilon^{-1} \ln(-p) + \left[1 - \ln(j\varepsilon) - \ln(-p) + \ln \left(1 - \frac{y}{q} \right) \right] j \right\} \\ + \varepsilon^{1/2} \sqrt{\frac{2}{\pi}} \frac{\sqrt{(1-y)y}}{y-q} \sin(N\pi y) \exp \left\{ \frac{y \ln(y) + (1-y) \ln(1-y) + \pi i}{\varepsilon} - j \ln(y-q) \right\}. \end{aligned} \quad (127)$$

Using (118) in (25) we have as $\varepsilon \rightarrow 0$, $0 < y < q$

$$\psi^+(y, z) \sim \ln(-p) + y \ln \left(\frac{q}{p} \right) - y\pi i + \left[1 - \ln(j\varepsilon) - \ln(-p) + \ln \left(1 - \frac{y}{q} \right) \right] j\varepsilon \quad (128)$$

$$\psi^-(y, z) \sim y \ln(y) + (1-y) \ln(1-y) + (1-y)\pi i - j \ln(y-q)\varepsilon \quad (129)$$

and from (26) we get

$$L^+(y, z) \sim \frac{1}{\sqrt{j\varepsilon}}, \quad L^-(y, z) \sim \sqrt{y(1-y)} \frac{1}{q-y} i. \quad (130)$$

Using (128)-(130) in (76) we have

$$\begin{aligned} K^{(7)}(y, z) \sim \cos \left(\frac{\pi y}{\varepsilon} \right) \frac{1}{\sqrt{2\pi j}} \exp \left\{ \frac{\ln(-p) + y \ln \left(\frac{q}{p} \right)}{\varepsilon} + \left[1 - \ln(j\varepsilon) - \ln(-p) + \ln \left(1 - \frac{y}{q} \right) \right] j \right\} \\ + \sqrt{y(1-y)} \frac{1}{y-q} \sin \left(\frac{\pi y}{\varepsilon} \right) \sqrt{\frac{2\varepsilon}{\pi}} \exp \left[\frac{y \ln(y) + (1-y) \ln(1-y) + \pi i}{\varepsilon} - j \ln(y-q) \right]. \end{aligned} \quad (131)$$

Matching (127) and 131 we obtain

$$A^{(11)}(y) = \exp \left[y \ln \left(\frac{q}{p} \right) \varepsilon^{-1} \right] \cos \left(\frac{\pi y}{\varepsilon} \right). \quad (132)$$

Hence, we conclude that

$$K_j^{(11)}(y) = \binom{N}{j} (-p)^{N-j} \left(\frac{q}{p} \right)^{Ny} \cos(N\pi y) \left(1 - \frac{y}{q} \right)^j + \binom{Ny}{N-j} \left(\frac{1-y}{q-y} \right)^{j+1} \quad (133)$$

for $z = 1 - O(\varepsilon)$ and $y \not\approx q$.

9.2 The corner layer at $(q, 1)$ (Region XII)

The approximation (133) ceases to be valid for $y \approx q$. Therefore, we need to find another expression, which holds in a neighborhood of the point $(q, 1)$. We introduce the new variable ξ defined by

$$y = q + \xi \sqrt{2pq\varepsilon}, \quad \xi = O(1). \quad (134)$$

Using (134) in (133) we have, as $\varepsilon \rightarrow 0$

$$\begin{aligned} K_j^{(11)}(y) &\sim \exp \left[\frac{p \ln(p) + q \ln(q)}{\varepsilon} + \xi \sqrt{\frac{2pq}{\varepsilon}} \ln \left(\frac{q}{p} \right) - \frac{j}{2} \ln(pq\varepsilon) \right] \\ &\times \left[\frac{(\sqrt{2}\xi)^j}{j!} \cos \left(\frac{p\pi}{\varepsilon} - \xi\pi \sqrt{\frac{2pq}{\varepsilon}} \right) - \sqrt{\frac{2}{\pi}} e^{\xi^2} (\sqrt{2}\xi)^{-j-1} \sin \left(\frac{p\pi}{\varepsilon} - \xi\pi \sqrt{\frac{2pq}{\varepsilon}} \right) \right]. \end{aligned} \quad (135)$$

Equation (135) suggests that we define a new function $R_j^{(12)}(\xi)$ by

$$K_n(x) = \exp \left[\frac{p \ln(p) + q \ln(q)}{\varepsilon} + \xi \sqrt{\frac{2pq}{\varepsilon}} \ln \left(\frac{q}{p} \right) - \frac{j}{2} \ln(pq\varepsilon) \right] R_j^{(12)}(\xi). \quad (136)$$

Using (136) in (4) we obtain, to leading order,

$$(j+1) R_{j+1}^{(12)} - \sqrt{2}\xi R_j + R_{j-1}^{(12)} = 0$$

which has the independent solutions [38]

$$R_j^{(12)}(\xi) = \frac{1}{j!} D_j(\sqrt{2}\xi), \quad \text{and} \quad R_j^{(12)}(\xi) = D_{-j-1}(\pm\sqrt{2}i\xi) (\pm i)^j.$$

Using (65) and matching with (135) we get

$$\begin{aligned} K_n(x) &\sim K_j^{(12)}(\xi) = \exp \left[\frac{p \ln(p) + q \ln(q)}{\varepsilon} + \xi \sqrt{\frac{2pq}{\varepsilon}} \ln \left(\frac{q}{p} \right) - \frac{j}{2} \ln(pq\varepsilon) \right] \exp \left(\frac{\xi^2}{2} \right) \\ &\times \left[\frac{1}{j!} D_j(\sqrt{2}\xi) \cos \left(\frac{p\pi}{\varepsilon} - \xi\pi \sqrt{\frac{2pq}{\varepsilon}} \right) - \frac{1}{\sqrt{2\pi}} \Lambda_j(\xi) \sin \left(\frac{p\pi}{\varepsilon} - \xi\pi \sqrt{\frac{2pq}{\varepsilon}} \right) \right], \end{aligned} \quad (137)$$

where the function $\Lambda_j : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$\Lambda_j(\xi) = i^{j+1} \left[D_{-j-1} \left(\sqrt{2}i\xi \right) + (-1)^{j+1} D_{-j-1} \left(-\sqrt{2}i\xi \right) \right]. \quad (138)$$

9.2.1 Matching the corner at $(q, 1)$ and the interior of $\mathbf{E}$

Finally, we verify the matching between (108) and (137). Using (118) and (134) in (25) we have,

$$\begin{aligned} \frac{\psi^+}{\varepsilon} &\sim \frac{q \ln(q) + p \ln(p) + p\pi i}{\varepsilon} + \xi \sqrt{2pq} \left[\ln \left(\frac{q}{p} \right) - \pi i \right] \varepsilon^{-1/2} \\ &\quad + \frac{j}{2} [1 - \ln(pqj\varepsilon) - \pi i] + \sqrt{2j}\xi i + \frac{\xi^2}{2} \\ \frac{\psi^-}{\varepsilon} &\sim \frac{q \ln(q) + p \ln(p) - p\pi i}{\varepsilon} + \xi \sqrt{2pq} \left[\ln \left(\frac{q}{p} \right) + \pi i \right] \varepsilon^{-1/2} \\ &\quad + \frac{j}{2} [1 - \ln(pqj\varepsilon) + \pi i] - \sqrt{2j}\xi i + \frac{\xi^2}{2}. \end{aligned}$$

Similarly, from (26) we get

$$L^+ \sim \frac{1}{\sqrt{2j\varepsilon}}, \quad L^- \sim \frac{1}{\sqrt{2j\varepsilon}}$$

and therefore

$$\begin{aligned} K^{(10)}(y, z) &\sim \frac{1}{\sqrt{\pi j}} \exp \left[\frac{q \ln(q) + p \ln(p)}{\varepsilon} + \xi \sqrt{2pq} \ln \left(\frac{q}{p} \right) \varepsilon^{-1/2} + \frac{\xi^2}{2} \right] \\ &\times \exp \left\{ \frac{j}{2} [1 - \ln(pqj\varepsilon)] \right\} \cos \left[\frac{p\pi}{\varepsilon} - \xi \sqrt{2pq}\pi \varepsilon^{-1/2} + \sqrt{2j}\xi - \frac{j\pi}{2} \right]. \end{aligned} \quad (139)$$

Using the formula [38]

$$D_n(\xi) \sim \sqrt{2} \exp \left\{ \frac{n}{2} [\ln(n) - 1] \right\} \cos \left[\sqrt{n}\xi - \frac{n\pi}{2} \right], \quad n \rightarrow \infty$$

in (137) yields

$$\begin{aligned} K_j^{(12)}(\xi) &\sim \exp \left[\frac{p \ln(p) + q \ln(q)}{\varepsilon} + \xi \sqrt{\frac{2pq}{\varepsilon}} \ln \left(\frac{q}{p} \right) - \frac{j}{2} \ln(pq\varepsilon) \right] \frac{1}{\sqrt{\pi j}} \exp \left\{ \frac{\xi^2}{2} + \frac{j}{2} [1 - \ln(j)] \right\} \\ &\times \left[\cos \left(\sqrt{j}\xi - \frac{j\pi}{2} \right) \cos \left(\frac{p\pi}{\varepsilon} - \xi \pi \sqrt{\frac{2pq}{\varepsilon}} \right) - \sin \left(\sqrt{j}\xi - \frac{j\pi}{2} \right) \sin \left(\frac{p\pi}{\varepsilon} - \xi \pi \sqrt{\frac{2pq}{\varepsilon}} \right) \right], \end{aligned} \quad (140)$$

where we have used

$$\Lambda_j(\xi) \sim \sqrt{\frac{2}{j}} \exp \left\{ \frac{j}{2} [1 - \ln(j)] \right\} \sin \left(\sqrt{j}\xi - \frac{j\pi}{2} \right).$$

Equations (139) and (140) are identical, after regrouping terms.




Figure 2: A sketch of the different asymptotic regions.

10 Summary and numerical results

Below we summarize our results for the various asymptotic approximations to $K_n(x)$ as $N \rightarrow \infty$, with

$$\varepsilon = \frac{1}{N}, \quad x = \frac{y}{\varepsilon}, \quad n = \frac{z}{\varepsilon} \quad 0 \leq y, z \leq 1.$$

(see Figure 2).

1. Region I: $n = O(1)$, $0 \leq y \leq 1$, $y \not\approx p$.

$$K_n(x) \sim K_n^{(1)}(y) = \frac{\varepsilon^{-n}}{n!} (y - p)^n.$$

(see Figure 3).




Figure 3: A comparison of $K_n(x)$ (solid curve) and $K_n^{(1)}(y)$ (ooo) for $n = 2$ with $\varepsilon = 0.01$ and $q = 0.64894783$.




Figure 4: A comparison of $K_n(x)$ (solid curve) and $K_n^{(2)}(\eta)$ (ooo) for $n = 2$ with $\varepsilon = 0.01$ and $q = 0.64894783$.

2. Region II: $n = O(1)$, $y \approx p$, $y = p + \eta\sqrt{2pq\varepsilon}$, $\eta = O(1)$.

$$K_n(x) \sim K_n^{(2)}(\eta) = \frac{\varepsilon^{-n/2}}{n!} \left(\frac{pq}{2}\right)^{n/2} H_n(\eta),$$

where $H_n(\eta)$ is the Hermite polynomial (see Figure 4).

3. Region III: $0 \leq y < Y^-(z)$, $0 < z < p$, where

$$Y^\pm(z) = p + (q - p)z \pm 2zU_0, \quad U_0(z) = \sqrt{\frac{pq(1-z)}{z}}.$$

$$K_n(x) \sim K^-(y, z) = \varepsilon^{1/2} \frac{1}{\sqrt{2\pi}} \exp[\varepsilon^{-1}\psi^-(y, z)] L^-(y, z),$$




Figure 5: A comparison of $K_n(x)$ (solid curve) and $K^-(y, z)$ (ooo) for $n = 10$ with $\varepsilon = 0.01$ and $q = 0.34894783$.

with

$$\psi^\pm(y, z) = \ln \left[(U^\pm)^{z-1} (U^\pm - p)^{1-y} (U^\pm + q)^y \right], \quad L^\pm(y, z) = \sqrt{\frac{(U^\pm - p)(U^\pm + q)}{z [(U^\pm)^2 - U_0^2]}}$$

and

$$U^\pm(y, z) = -\frac{1}{2} \left(\frac{p-y}{z} + q - p \right) \pm \frac{1}{2} \sqrt{\left(\frac{p-y}{z} + q - p \right)^2 - 4(U_0)^2}$$

(see Figure 5).

4. Region IV: $Y^+(z) < y \leq 1$, $0 < z < q$.

$$K_n(x) \sim K^+(y, z) = \varepsilon^{1/2} \frac{1}{\sqrt{2\pi}} \exp [\varepsilon^{-1} \psi^+(y, z)] L^+(y, z).$$




Figure 6: A comparison of $K_n(x)$ (solid curve) and $K^+(y, z)$ (ooo) for $n = 10$ with $\varepsilon = 0.01$ and $q = 0.34894783$.

(see Figure 6).

5. Region V: $x = O(1)$, $p < z < 1$.

$$K_n(x) \sim K^{(5)}(x, z) = \frac{\varepsilon^{1/2}}{\sqrt{2\pi}\sqrt{z(1-z)}} \cos(\pi x) \left(\frac{z-p}{p} \right)^x \exp \left[\frac{\phi_0(z)}{\varepsilon} \right] \\ - \frac{\varepsilon}{\pi} \frac{x}{z-p} \Gamma(x) \sin(\pi x) \left(\frac{q\varepsilon}{z-p} \right)^x \exp \left[\frac{(z-1)\ln(q) + \pi iz}{\varepsilon} \right]$$

where

$$\phi_0(z) = (z-1)\ln(1-z) - z\ln(z) + z\ln(-p)$$

and $\Gamma(x)$ is the Gamma function (see Figure 7).




Figure 7: A comparison of $K_n(x)$ (solid curve) and $K^{(5)}(x, z)$ (ooo) for $n = 80$ with $\varepsilon = 0.01$ and $q = 0.74894783$.




Figure 8: A comparison of $K_n(x)$ (solid curve) and $K^{(6)}(x, u)$ (ooo) for $n = 25$ with $\varepsilon = 0.01$ and $q = 0.74894783$. Note that with this choice of parameters, $u = 0.024265$.

6. Region VI: $x = O(1)$, $z \approx p$, $z = p - u\sqrt{pq\varepsilon}$, $u = O(1)$.

$$K_n(x) \sim K^{(6)}(x, u) = \frac{\varepsilon^{1/2}}{\sqrt{2\pi pq}} \left[\sqrt{\frac{q\varepsilon}{p}} \right]^x D_x(u) \\ \times \exp \left[\frac{\pi i p - q \ln(q)}{\varepsilon} + \frac{u\sqrt{pq}\pi i - u\sqrt{pq} \ln(q)}{\sqrt{\varepsilon}} - \frac{u^2}{4} \right],$$

where $D_x(u)$ is the parabolic cylinder function (see Figure 8).

7. Region VII: $0 \ll y < Y^-(z)$, $p < z < 1$.

$$K_n(x) \sim K^{(7)}(y, z) = \exp \left(\frac{\pi i y}{\varepsilon} \right) \left[\cos \left(\frac{\pi y}{\varepsilon} \right) K^+(y, z) + 2i \sin \left(\frac{\pi y}{\varepsilon} \right) K^-(y, z) \right]$$

(see Figure 9).




Figure 9: A comparison of $K_n(x)$ (solid curve) and $K^{(7)}(y, z)$ (ooo) for $n = 35$ with $\varepsilon = 0.025$ and $q = 0.74894783$.




Figure 10: A comparison of $K_n(x)$ (solid curve) and $K^{(8)}(\beta, z)$ (ooo) for $n = 10$ with $\varepsilon = 0.01$ and $q = 0.34894783$.

8. Region VIII: $y \approx Y^-(z)$, $0 < z < p$, $y = Y^-(z) - \beta\varepsilon^{2/3}$, $\beta = O(1)$.

$$K_n(x) \sim K^{(8)}(\beta, z) = \varepsilon^{1/3} \exp \left[\varepsilon^{-1} \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{-1/3} \right] \text{Ai} \left[\Theta^{2/3} \beta \right] \frac{\Theta^{-1/3}}{\sqrt{z U_0}},$$

where

$$\psi_0(z) = z\pi i + (z - 1) \ln(U_0) + Y^-(z) \ln(U_0 - q) + [1 - Y^-(z)] \ln(U_0 + p),$$

$$\Theta(z) = \sqrt{\frac{U_0}{z}} \frac{1}{(U_0 + p)(U_0 - q)}$$

and $\text{Ai}(\cdot)$ is the Airy function (see Figure 10).

9. Region IX: $y \approx Y^-(z)$, $p < z < 1$.

$$K_n(x) \sim K^{(9)}(\beta, z) = \varepsilon^{1/3} \exp \left[\varepsilon^{-1} \psi_0(z) + \ln \left(\frac{U_0 + p}{U_0 - q} \right) \beta \varepsilon^{-1/3} \right] \\ \times \frac{1}{2} \frac{\vartheta^{-1/3}}{\sqrt{zU_0}} \left[\lambda^+(\beta, z) \text{Ai}(\vartheta^{2/3} \beta) + i \lambda^-(\beta, z) \text{Bi}(\vartheta^{2/3} \beta) \right],$$

where $\vartheta(z) = -\Theta(z)$,

$$\lambda^\pm(\beta, z) = \exp \left\{ \frac{2\pi i [Y^-(z) - \beta \varepsilon^{2/3}]}{\varepsilon} \right\} \pm 1.$$

and $\text{Ai}(\cdot)$, $\text{Bi}(\cdot)$ are the Airy functions (see Figure 11).

10. Region X: $Y^-(z) < y < Y^+(z)$, $0 < z < 1$.

$$K_n(x) \sim K^{(10)}(y, z) = K^+(y, z) + K^-(y, z)$$

(see Figure 12).

11. Region XI: $n \approx N$, $n = N - j$, $j \in \mathbb{Z}$, $0 < y < 1$, $y \not\approx q$.

$$K_n(x) \sim K_j^{(11)}(y) = \binom{N}{j} (-p)^{N-j} \left(\frac{q}{p} \right)^{Ny} \cos(N\pi y) \left(1 - \frac{y}{q} \right)^j + \binom{Ny}{N-j} \left(\frac{1-y}{q-y} \right)^{j+1}$$

(see Figure 13).

12. Region XII: $n \approx N$, $y \approx q$, $y = q + \xi \sqrt{2pq\varepsilon}$, $\xi = O(1)$.

$$K_n(x) \sim K_j^{(12)}(\xi) = \exp \left[\frac{p \ln(p) + q \ln(q)}{\varepsilon} + \xi \sqrt{\frac{2pq}{\varepsilon}} \ln \left(\frac{q}{p} \right) - \frac{j}{2} \ln(pq\varepsilon) \right] \exp \left(\frac{\xi^2}{2} \right) \\ \times \left[\frac{1}{j!} D_j \left(\sqrt{2\xi} \right) \cos \left(\frac{p\pi}{\varepsilon} - \xi \pi \sqrt{\frac{2pq}{\varepsilon}} \right) - \frac{1}{\sqrt{2\pi}} \Lambda_j(\xi) \sin \left(\frac{p\pi}{\varepsilon} - \xi \pi \sqrt{\frac{2pq}{\varepsilon}} \right) \right],$$

where the function $\Lambda_j : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$\Lambda_j(\xi) = i^{j+1} \left[D_{-j-1} \left(\sqrt{2i\xi} \right) + (-1)^{j+1} D_{-j-1} \left(-\sqrt{2i\xi} \right) \right]$$

(see Figure 14).




Figure 11: A comparison of $K_n(x)$ (solid curve) and $K^{(9)}(\beta, z)$ (ooo) for $n = 40$ with $\varepsilon = 0.02$ and $q = 0.74894783$.




Figure 12: A comparison of $K_n(x)$ (solid curve) and $K^{(10)}(y, z)$ (ooo) for $n = 40$ with $\varepsilon = 0.02$ and $q = 0.74894783$.




Figure 13: A comparison of $K_n(x)$ (solid curve) and $K_j^{(11)}(y)$ (ooo) for $n = 19$ with $\varepsilon = 0.05$ and $q = 0.74894783$.




Figure 14: A comparison of $K_n(x)$ (solid curve) and $K_j^{(12)}(\xi)$ (ooo) for $n = 20$ with $\varepsilon = 0.05$ and $q = 0.74894783$.

References

- [1] M. Abramowitz and I. A. Stegun, editors. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. Dover Publications Inc., New York, 1992. Reprint of the 1972 edition.
- [2] N. M. Atakishiyev, G. S. Pogosyan, L. E. Vicent, and K. B. Wolf. Separation of discrete variables in the 2-dim finite oscillator. In *Quantum theory and symmetries (Kraków, 2001)*, pages 255–260. World Sci. Publishing, River Edge, NJ, 2002.
- [3] L. A. Bassalygo. Generalization of Lloyd’s theorem to arbitrary alphabet. *Problems of Control and Information Theory/Problemy Upravleniya i Teorii Informacii*, 2(2):133–137, 1973.
- [4] L. Chihara and D. Stanton. Zeros of generalized Krawtchouk polynomials. *J. Approx. Theory*, 60(1):43–57, 1990.
- [5] P. Delsarte. An algebraic approach to the association schemes of coding theory. *Philips Res. Rep. Suppl.*, (10):vi+97, 1973.
- [6] D. Dominici and C. Knessl. Asymptotic analysis by the saddle point method of a data-handling system with many sources. *The Journal of Applied Mathematics and Stochastic Analysis*. To appear.
- [7] P. D. Dragnev and E. B. Saff. A problem in potential theory and zero asymptotics of Krawtchouk polynomials. *J. Approx. Theory*, 102(1):120–140, 2000.
- [8] P. Feinsilver and R. Schott. Krawtchouk polynomials and finite probability theory. In *Probability measures on groups, X (Oberwolfach, 1990)*, pages 129–135. Plenum, New York, 1991.
- [9] L. Habsieger. Integer zeros of q -Krawtchouk polynomials in classical combinatorics. *Adv. in Appl. Math.*, 27(2-3):427–437, 2001. Special issue in honor of Dominique Foata’s 65th birthday (Philadelphia, PA, 2000).
- [10] L. Habsieger. Integral zeroes of Krawtchouk polynomials. In *Codes and association schemes (Piscataway, NJ, 1999)*, volume 56 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 151–165. Amer. Math. Soc., Providence, RI, 2001.
- [11] L. Habsieger and D. Stanton. More zeros of Krawtchouk polynomials. *Graphs Combin.*, 9(2):163–172, 1993.
- [12] M. E. H. Ismail and P. Simeonov. Strong asymptotics for Krawtchouk polynomials. *J. Comput. Appl. Math.*, 100(2):121–144, 1998.

- [13] C. Ivan. A multidimensional nonlinear growth, birth and death, emigration and immigration process. In *Proceedings of the Fourth Conference on Probability Theory (Braşov, 1971)*, pages 421–427. Editura Acad. R. S. R., Bucharest, 1973.
- [14] R. Koekoek and R. F. Swarttouw. The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue. Technical Report 98-17, Delft University of Technology, 1998. <http://aw.twi.tudelft.nl/~koekoek/askey/>.
- [15] I. Krasikov. Bounds for the Christoffel-Darboux kernel of the binary Krawtchouk polynomials. In *Codes and association schemes (Piscataway, NJ, 1999)*, volume 56 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 193–198. Amer. Math. Soc., Providence, RI, 2001.
- [16] I. Krasikov and S. Litsyn. On integral zeros of Krawtchouk polynomials. *J. Combin. Theory Ser. A*, 74(1):71–99, 1996.
- [17] I. Krasikov and S. Litsyn. On the distance distributions of BCH codes and their duals. *Des. Codes Cryptogr.*, 23(2):223–231, 2001.
- [18] I. Krasikov and S. Litsyn. Survey of binary Krawtchouk polynomials. In *Codes and association schemes (Piscataway, NJ, 1999)*, volume 56 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 199–211. Amer. Math. Soc., Providence, RI, 2001.
- [19] N. N. Lebedev. *Special functions and their applications*. Dover Publications Inc., New York, 1972. Translated from the Russian.
- [20] H. W. Lenstra, Jr. Two theorems on perfect codes. *Discrete Math.*, 3:125–132, 1972.
- [21] V. I. Levenshtein. Krawtchouk polynomials and universal bounds for codes and designs in Hamming spaces. *IEEE Trans. Inform. Theory*, 41(5):1303–1321, 1995.
- [22] X.-C. Li and R. Wong. A uniform asymptotic expansion for Krawtchouk polynomials. *J. Approx. Theory*, 106(1):155–184, 2000.
- [23] S. P. Lloyd. Binary block coding. *Bell System Tech. J.*, 36:517–535, 1957.
- [24] M. Lorente. Quantum mechanics on discrete space and time. In *New developments on fundamental problems in quantum physics (Oviedo, 1996)*, volume 81 of *Fund. Theories Phys.*, pages 213–224. Kluwer Acad. Publ., Dordrecht, 1997.
- [25] F. J. MacWilliams and N. J. A. Sloane. *The theory of error-correcting codes. I*. North-Holland Publishing Co., Amsterdam, 1977. North-Holland Mathematical Library, Vol. 16.

- [26] P. Nevai. Two of my favorite ways of obtaining asymptotics for orthogonal polynomials. In *Anniversary volume on approximation theory and functional analysis (Oberwolfach, 1983)*, volume 65 of *Internat. Schriftenreihe Numer. Math.*, pages 417–436. Birkhäuser, Basel, 1984.
- [27] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov. *Classical orthogonal polynomials of a discrete variable*. Springer Series in Computational Physics. Springer-Verlag, Berlin, 1991. Translated from the Russian.
- [28] F. W. J. Olver. *Asymptotics and special functions*. AKP Classics. A K Peters Ltd., Wellesley, MA, 1997. Reprint of the 1974 original.
- [29] A. Poli and L. Huguet. *Error correcting codes*. Prentice Hall International, Hemel Hempstead, 1992. Translated from the 1989 French original.
- [30] W.-Y. Qiu and R. Wong. Asymptotic expansion of the Krawtchouk polynomials and their zeros. *Comput. Methods Funct. Theory*, 4(1):189–226, 2004.
- [31] W. Schoutens. *Stochastic processes and orthogonal polynomials*, volume 146 of *Lecture Notes in Statistics*. Springer-Verlag, New York, 2000.
- [32] I. I. Sharapudinov. Asymptotic properties of Krawtchouk polynomials. *Mat. Zametki*, 44(5):682–693, 703, 1988.
- [33] N. J. A. Sloane. An introduction to association schemes and coding theory. In *Theory and application of special functions (Proc. Advanced Sem., Math. Res. Center, Univ. Wisconsin, Madison, Wis., 1975)*, pages 225–260. Math. Res. Center, Univ. Wisconsin, Publ. No. 35. Academic Press, New York, 1975.
- [34] P. Solé. An inversion formula for Krawtchouk polynomials with applications to coding theory. *J. Inform. Optim. Sci.*, 11(2):207–213, 1990.
- [35] R. J. Stroeker and B. M. M. de Weger. On integral zeroes of binary Krawtchouk polynomials. *Nieuw Arch. Wisk. (4)*, 17(2):175–186, 1999.
- [36] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, R.I., fourth edition, 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.
- [37] Z. Wang and R. Wong. Asymptotic expansions for second-order linear difference equations with a turning point. *Numer. Math.*, 94(1):147–194, 2003.
- [38] E. T. Whittaker and G. N. Watson. *A course of modern analysis*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1996. Reprint of the fourth (1927) edition.

- [39] A. Zhedanov. Oscillator $9j$ -symbols, multidimensional factorization method, and multivariable Krawtchouk polynomials. In *Calogero-Moser-Sutherland models (Montréal, QC, 1997)*, CRM Ser. Math. Phys., pages 549–561. Springer, New York, 2000.